



Lessons from the 2018–2019 European droughts: a collective need for unifying drought risk management

Veit Blauhut¹, Michael Stoelzle¹, Lauri Ahopelto^{2,3}, Manuela I. Brunner¹, Claudia Teutschbein⁴, Doris E. Wendt^{5,6}, Vytautas Akstinas⁷, Sigrid J. Bakke⁸, Lucy J. Barker⁹, Lenka Bartošová¹⁰, Agrita Briede¹¹, Carmelo Cammalleri¹², Ksenija Cindrić Kalin¹³, Lucia De Stefano¹⁴, Miriam Fendeková¹⁵, David C. Finger^{16,17}, Marijke Huysmans¹⁸, Mirjana Ivanov¹⁹, Jaak Jaagus²⁰, Jiří Jakubínský¹⁰, Svitlana Krakovska²¹, Gregor Laaha²², Monika Lakatos²³, Kiril Manevski²⁴, Mathias Neumann Andersen²⁴, Nina Nikolova²⁵, Marzena Osuch²⁶, Pieter van Oel²⁷, Kalina Radeva²⁵, Renata J. Romanowicz²⁶, Elena Toth²⁸, Mirek Trnka¹⁰, Marko Urošev²⁹, Julia Urquijo Reguera³⁰, Eric Sauquet³¹, Aleksandra Stevkov³², Lena M. Tallaksen⁸, Iryna Trofimova²¹, Anne F. Van Loon³³, Michelle T. H. van Vliet³⁴, Jean-Philippe Vidal³¹, Niko Wanders³⁴, Micha Werner³⁵, Patrick Willems³⁶, and Nenad Živković³⁷

¹Environmental Hydrological Systems, Faculty of Environment and Natural Resources, University of Freiburg, Freiburg, Germany

²Water and Development Research Group, School of Engineering, Aalto University, Aalto, Finland

³Freshwater Centre, Finnish Environment Institute, Helsinki, Finland

⁴Program for Air, Water and Landscape Sciences – Hydrology, Department of Earth Sciences, Uppsala University, Uppsala, Sweden

⁵School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham, UK

⁶Department of Civil Engineering, University of Bristol, Bristol, UK

⁷Laboratory of Hydrology, Lithuanian Energy Institute, Kaunas, Lithuania

⁸Department of Geosciences, University of Oslo, Oslo, Norway

⁹UK Centre for Ecology & Hydrology, Wallingford, UK

¹⁰Global Change Research Institute CAS, Brno, Czech Republic

¹¹Faculty of Geography and Earth Sciences, University of Latvia, Riga, Latvia

¹²Joint Research Centre, European Commission, Ispra, Italy

¹³Division for the Development of Climate Products and Applications, Croatian Meteorological and Hydrological Service, Zagreb, Croatia

¹⁴Facultad de Ciencias Geológicas, Universidad Complutense de Madrid, Madrid, Spain

¹⁵Department of Hydrogeology, Faculty of Natural Sciences, Comenius University in Bratislava, Mlynska dolina, Ilkovičova 6, 842 15 Bratislava 4, Slovakia

¹⁶School of engineering, Reykjavik University, Reykjavik, Iceland

¹⁷Energieinstitut an der Johannes Kepler Universität, Linz, Austria

¹⁸Department of Hydrology and Hydraulic Engineering, Vrije Universiteit Brussel, Brussels, Belgium

¹⁹Institute of Hydrometeorology and Seismology, Podgorica, Montenegro

²⁰Department of Geography, Institute of Ecology and Earth Sciences, University of Tartu, Tartu, Estonia

²¹Laboratory of Applied Climatology, Ukrainian Hydrometeorological Institute, Kyiv, Ukraine

²²Institute of Statistics, University of Natural Resources and Life Sciences, Vienna, Austria

²³Hungarian Meteorological Service, Budapest, Hungary

²⁴Department of Agroecology, Aarhus University, Tjele, Denmark

²⁵Department of Climatology, Hydrology and Geomorphology, Faculty of Geology and Geography, Sofia University “St. Kliment Ohridski”, Sofia, Bulgaria

²⁶Institute of Geophysics, Polish Academy of Sciences, Warsaw, Poland

²⁷Water Resources Management Group, Wageningen University, Wageningen, the Netherlands

²⁸Department of Civil, Chemical, Environmental and Materials Engineering, University of Bologna, Bologna, Italy

²⁹Geographical Institute “Jovan Cvijić”, Serbian Academy of Sciences and Arts, Belgrade, Serbia

³⁰Department of Agroforestry Engineering, Escuela Técnica Superior de Ingeniería Agronómica y de Biosistemas, Universidad Politécnica de Madrid, Madrid, Spain

³¹RiverLy, INRAE, Villeurbanne, France

³²Department of Meteorology, National Hydrometeorological Service, Skopje, North Macedonia

³³Institute for Environmental Studies, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands

³⁴Department of Physical Geography, Utrecht University, Utrecht, the Netherlands

³⁵Water Resources & Ecosystems Department, IHE Delft Institute for Water Education, Delft, the Netherlands

³⁶Hydraulics and Geotechnics Section, Department of Civil Engineering, KU Leuven, Leuven, Belgium

³⁷Faculty of Geography, University of Belgrade, Belgrade, Serbia

Correspondence: Veit Blauhut (veit.blauhut@hydrology.uni-freiburg.de)

Received: 24 September 2021 – Discussion started: 13 October 2021

Revised: 29 April 2022 – Accepted: 9 May 2022 – Published: 29 June 2022

Abstract. Drought events and their impacts vary spatially and temporally due to diverse pedo-climatic and hydrologic conditions, as well as variations in exposure and vulnerability, such as demographics and response actions. While hazard severity and frequency of past drought events have been studied in detail, little is known about the effect of drought management strategies on the actual impacts and how the hazard is perceived by relevant stakeholders. In a continental study, we characterised and assessed the impacts and the perceptions of two recent drought events (2018 and 2019) in Europe and examined the relationship between management strategies and drought perception, hazard, and impact. The study was based on a pan-European survey involving national representatives from 28 countries and relevant stakeholders responding to a standard questionnaire. The survey focused on collecting information on stakeholders’ perceptions of drought, impacts on water resources and beyond, water availability, and current drought management strategies on national and regional scales. The survey results were compared with the actual drought hazard information registered by the European Drought Observatory (EDO) for 2018 and 2019. The results highlighted high diversity in drought perception across different countries and in values of the implemented drought management strategies to alleviate impacts by increasing national and sub-national awareness and resilience. The study identifies an urgent need to further reduce drought impacts by constructing and implementing a European macro-level drought governance approach, such as a directive, which would strengthen national drought management and mitigate damage to human and natural assets.

1 Introduction

1.1 Drought impacts in Europe

During recent decades, different parts of Europe have been affected by several severe, large-scale drought events, e.g. in 2003, 2007, 2011, 2012, 2015, 2017, 2018, 2019, and 2020 (Baruth et al., 2020; Boergens et al., 2020; Cindrić Kalin et al., 2016; García-Herrera et al., 2019; Hänsel et al., 2019; Ionita et al., 2017; Laaha et al., 2017; McKee et al., 1993). Each of these droughts was unique in terms of severity, spatio-temporal extent, and associated direct and indirect impacts on human and natural resources (Stahl et al., 2016). Cammalleri et al. (2020) estimated drought-related losses in the European Union (EU) to be about EUR 9 billion annually. The largest share of these losses is typically seen in agricultural, energy, and public water supply sectors (Cammalleri et al., 2020), triggered mainly by agricultural (soil moisture deficit) and hydrological drought (deficit in river flow and groundwater; Van Lanen et al., 2016). These sectoral losses likely represent only part of the actual drought impacts as indirect, intangible, or subtle impacts are more difficult to identify and quantify, such as adverse effects on ecosystem services and human health (Erian et al., 2021). According to the European Drought Impact report Inventory (EDII; Stahl et al., 2016), further impacts on aquaculture, ecosystems, and human and public safety, as well as conflicts between sectoral water users, have been reported. Herein, the occurrence and the composition of drought impacts are assumed to greatly vary with regional and national exposure, perception, and vulnerability to droughts (e.g. Stahl et al., 2016).

1.2 Drought management in Europe

A key element to mitigate drought impact is to respond promptly, i.e. implement drought management planning strategies and associated action plans (UNDRR, 2019). However, a directive for drought risk management does not exist

on a near-continental scale, such as in the European Union (EU; Hervás-Gómez et al., 2019), despite the identified potential for reducing emergency management costs through proactive management (Cammalleri et al., 2020; Howarth, 2018). So far, “droughts have only been succinctly dealt with in the Water Framework Directive with no compulsory actions” (Hervás-Gómez et al., 2019). However, recommendations are not adopted in all relevant/major river basin districts (European Commission, 2019). The “European Commission’s Communication on water scarcity and drought” and the “Blueprint to Safeguard Europe’s Water Resources” (European Commission, 2012) directly tackle drought and address current flaws and policy gaps. These documents have received a mixed response, ranging from “the Communication is still weak and lacks teeth in the policy landscape” (Stein et al., 2016) to “it is hoped to lead to an EU water policy development in a long term” (Hervás-Gómez et al., 2019). However, some countries historically have been, and are, more prone to drought compared to others due to their pedo-climatic settings, and although drought risk management does exist in these countries through national legislation, it mostly happens indirectly via policy-making regarding environmental protection, soil management, or water and climate adaptation (e.g. Zoth et al., 2019; Hanger-Kopp and Palka, 2020). Moreover, a number of technical guidelines exist to support the development and the implementation of national drought resilience, adaptation, and management plans (e.g. UNCCD, 2019). In fact, different national legal approaches not being internationally coordinated can create conflicts, i.e. water scarcity in one region/country at the cost of another, such as the case of the Blue Nile between Egypt and Ethiopia (Bastawesy, 2015) or the Danube between Hungary and Slovakia (Vuković et al., 2014). Therefore, a coordinated approach is required. Trnka et al. (2018) suggested to improve the understanding of triggers causing paradigm shifts from response-based to proactive drought management and policies as a priority research question.

1.3 The 2018 and 2019 European droughts

For several successive years, large parts of Europe were affected by severe and widespread droughts, which highlighted the vulnerability of its socio-economic and environmental systems. The 2018 event was special because of both rainfall deficits and high temperatures in many European countries (Rosner et al., 2019), with record-breaking high temperatures in several regions (Bakke et al., 2020), which reached otherwise cool and humid northern regions. This compound hot-dry event led to major impacts in north-central and north-eastern Europe, particularly affecting agriculture, livestock farming, and forestry (Bakke et al., 2020; Beillouin et al., 2020; Rosner et al., 2019; Salmoral et al., 2020; Schuldt et al., 2020; Thompson et al., 2020) as reported for Sweden, Finland, Estonia, Lithuania, Latvia, Denmark, the Netherlands, Belgium, Germany, the United Kingdom (UK), and

eastern France (Moravec et al., 2021; Turner et al., 2021). The propagation of the meteorological drought resulted in low reservoir levels and river discharge, which impaired public water supply, leading to partial shut downs of nuclear power plants and triggering massive fish deaths in upstream watersheds (e.g. de Brito, 2021). In contrast to central and northern Europe, the western Mediterranean countries experienced above-average wet conditions in 2018 after having experienced a very severe drought on the Iberian Peninsula in 2016–2017 and in Italy in 2017 (García-Herrera et al., 2019; Rita et al., 2020), while the eastern Mediterranean experienced below-average dry conditions (DriDanube-Watch, 2018). In contrast to 2018, the 2019 drought was centred on eastern Germany, the Czech Republic, and Poland before spreading westward (Boergens et al., 2020). The most affected regions were still suffering from large water balance deficits from the 2018 drought (Boergens et al., 2020) at the start of 2019. Hari et al. (2020) declared the period 2018–2019 in central Europe a 2-year drought event unprecedented in severity in the last 250 years, whereas Büntgen et al. (2021) show an accumulation of drought signals in central Europe over five summers, i.e. 2014–2018.

1.4 Drought risk and perception

The hydro-climatic aspects of past drought events have been studied in detail (e.g. Barker et al., 2019; Hisdal and Tallaksen, 2003; Dai, 2013; Cheval et al., 2014; Jaagus et al., 2021; Laaha et al., 2017; Radeva et al., 2018; Spinoni et al., 2015, 2018), whereas knowledge of the relationship between drought management, perception, and impacts remains limited (Blauhut, 2020; Hagenlocher et al., 2019; Kreibich et al., 2019). Understanding how different stakeholders perceive a specific drought event and its potential impacts can contribute to defining and successfully implementing drought mitigation measures adapted to a site-specific context (Alduce et al., 2017). Only a few studies have analysed relationships between drought perceptions and impacts. For instance, Teutschbein et al. (2019) assessed the link between perceived drought severity, impacts, preparedness, and management and measured hydrological drought impacts for two consecutive drought events (2017 and 2018) in Sweden. Although the authors did not find a significant relationship between the perceived level of drought impacts and the presence of a drought action plan, there was evidence that regions with a drought action plan applied significantly more measures in their drought response. Furthermore, the perceived drought severity in these regions did not match the observed severity of meteorological and hydrological droughts in Sweden: decision makers consistently overestimated the severity of mild drought events, while they underestimated more extreme drought conditions. In contrast, Blauhut et al. (2016) identified “drought awareness” and “drought management plans” as vulnerability factors driving drought risk for certain impact categories, such as agriculture and livestock farming,

public water supply, and freshwater ecosystems. The analysis of Blauhut et al. (2015) suggested that while national and international water management policies and guidelines may have decreased vulnerability, they may also have increased awareness and recognition of environmental impacts, leading to an increased number of reported drought impacts. Hence, previous statements on the relationship between the existence of drought risk management plans and drought impacts cannot be generalised.

1.5 Study aim

The aim of this paper is to assess how monitored drought hazard severity relates to drought perception and drought management strategies. We hypothesise that perceived drought impacts are not necessarily related to the severity of the drought hazard but are strongly influenced by national awareness and drought management strategies. To verify this hypothesis, we investigated how the droughts of 2018 and 2019 in 28 European countries were related to (a) the drought hazard as monitored by the European Drought Observatory (EDO), (b) drought management actions taken in the different countries, (c) drought perception by water managers and agencies, and (d) drought awareness. National drought perceptions, management, and impacts were studied using a pan-European survey. On the basis of this survey, we discuss the potential benefits of a European drought directive, similar to the Floods Directive (2007/60/EC) with respect to reducing drought vulnerability and impacts by macro-level governance.

2 Data

In order to evaluate the hypothesis, two different types of spatial data were collected and compared: (i) drought information as monitored by the EDO (<https://edo.jrc.ec.europa.eu>, last access: 1 April 2022) and (ii) information on drought impacts, perception, and state of drought management plans collected through a pan-European survey targeting water managers and water agencies. Note that Kosovo was not investigated disaggregated from Serbia (please see “Disclaimer”).

The hydro-climatic situation in 2018 and 2019 was described using a set of drought indices compiled by EDO for a variety of drought types including meteorological drought (Standardised Precipitation Index (SPI) for accumulation periods of 1, 3, 6, 9, and 12 months), soil moisture drought (Soil Moisture Anomaly; SM), hydrological drought (Low Flow Index, LFI, representing the discharge anomaly with respect to a daily threshold), and vegetation drought (anomaly of Fraction of Absorbed Photosynthetically Active Radiation; FAPAR). The SPI is given at a monthly resolution, whereas the other indices are presented in 10 d non-overlapping intervals. To increase comparability of the four indices, the EDO

Table 1. Drought indices and their associated drought classes. SPI, FAPAR, SM, and LFI are, respectively, Standardised Precipitation Index, Fraction of Accumulated Photosynthetically Active Radiation, Soil Moisture, and Low Flow Index.

Indices	No drought	Moderate drought	Severe drought	Extreme drought
SPI, FAPAR, SM	> -1	-1 to -1.5	-1.5 to -2	< -2
LFI	0–0.25	0.25–0.5	0.5–0.75	0.75–1

data were further classified into categorical drought classes: no drought, moderate drought, severe drought, and extreme drought. The standardised products SPI, FAPAR, and SM are categorised following McKee et al. (1993) (Table 1), and the LFI is computed from the daily streamflow values produced by the LISFLOOD hydrological model. The drought classification scheme used for LFI is taken from the European Drought Observatory (Table 1). These drought classes are in operational use at the EDO. Furthermore, the FAPAR was restricted to the warm season in Europe from April to August and was not monitored for Iceland. Detailed information on the drought indices and drought classes applied herein can be found in the corresponding EDO indicator fact sheets (<https://edo.jrc.ec.europa.eu/edov2/php/index.php?id=1101>, last access: 1 April 2022).

In order to assess the country-specific perception of drought, management, and impacts with a focus on 2018 and 2019, a pan-European survey was designed by the International Association of Hydrological Science (IAHS) – Panta Rhei “Drought in the Anthropocene” working group. National representatives of each country were selected and assigned responsibility to translate, distribute, and evaluate the survey and all associated communication and feedback. The network of national representatives developed out of our active Panta Rhei “Drought in the Anthropocene” group but also partly from the Euro-FRIEND “Low Flow and Drought” group. The idea was to have representatives affiliated with science or governmental agencies. In doing so, we expected a neutral point of view and comprehensive knowledge on the different aspects we were interested in. Furthermore, we expected such persons to be well networked and thus constitute a representative sample of stakeholders within each country. The survey targeted representatives of water management organisations and water agencies. Survey respondents were selected by the national representatives aiming to provide a balanced view of national opinions and drought management practices (or actions), as well as local and regional knowledge within each country. The content of the survey was adapted from Teutschbein et al. (2019), who studied 290 Swedish municipalities to evaluate the relationship between perceived drought severity, impact, preparedness, and management, aiming to compare stakeholder perception with hydrological drought indices. Note the perception of heat was not investigated.

The 26 questions of the survey covered the following themes:

- respondent background and the available water resource(s) used/managed,
- general perception of drought and associated risks,
- drought risk-related concepts and the drought management applied, and
- perception and impacts of the 2018 and the 2019 drought events.

The survey questions can be found in Table S1 in the Supplement. The paper and the figures displayed in the main body present a synthesis and insights from the pan-European comparison of the responses. More detailed aspects of the individual country responses are shown in Figs. S1–S7 of the Supplement.

3 Results and discussion

3.1 The drought events of 2018 and 2019 – hydro-meteorological results

The drought indices of the pan-European droughts in 2018 and 2019 are presented in Fig. 1a, which shows the meteorological drought conditions for selected months and the percentage area of land under severe or extreme drought. National results at a monthly resolution are shown in Fig. S1. Overall, the 2018 meteorological drought (as defined by SPI-3, SPI-6, SPI-9, and SPI-12) affected mainly central and northern Europe. The Benelux countries, Germany, Denmark, Sweden, and Finland showed an especially high spatial coverage of severe or extreme drought hazard. In early spring, rainfall deficits started in the north, i.e. Norway, Sweden, Finland, Lithuania, and Latvia (Fig. S1), and accumulated over central and northern Europe, peaking in the summer with high shares of extreme drought hazard at short accumulation periods (SPI-1). Strong soil moisture deficits in the summer were detected in regions affected by strong precipitation deficits over multiple months (SPI-3, SPI-6; Fig. S1). At the European scale, soil moisture deficits were especially high in northern Europe from June to August, and the area under severe drought in central Europe peaked in October and November (Fig. 1).

The hydrological drought of 2018 followed a similar spatial pattern as the meteorological drought (Fig. 1b), with severe hazard levels in the Benelux countries, Germany, the Czech Republic, Norway, and Sweden. The maximum spatial coverage of severe or extreme low flows in northern Europe occurred in June and July, in particular for countries where rainfall deficits continued and more intense deficits developed (SPI-9, SPI-12; Fig. S1). The maximum coverage of the 2018 hydrological drought in central Europe occurred in October and November (Fig. 1b).

Vegetation drought indicated by FAPAR was the most severe in Denmark and was contrasting with the drought signals of the other indices. For example, large parts of Belarus and France were under severe or extreme drought, while only small parts of Sweden and Finland were affected. In countries where precipitation deficits continued to accumulate over the 2018–2019 winter period, water deficits resulted in country-specific low flow conditions (Fig. 1b). The multi-year drought 2018–2019 particularly affected Belgium, Belarus, the Czech Republic, Germany, Finland, Latvia, Luxembourg, Lithuania, the Netherlands, Poland, Sweden, Switzerland, and the UK. Furthermore, Iceland also experienced an exceptionally long dry period in 2019. However, the effects were not as intense as in mainland Europe, mainly due to the numerous ice caps that provided ice melt, an extensive snow cover during winter (Helmert et al., 2018), a subpolar climate, and warm and humid ocean winds that could generate local rain events (Finger, 2018).

The 2019 drought was overall less severe compared to 2018, except for Iceland, which experienced an unusually long dry period. The centre of the meteorological drought moved eastwards with large areas under severe or extreme drought in Lithuania, Belarus, and Ukraine. Despite the lower intensity of the 2019 meteorological drought, soil moisture deficits remained high in central Europe, especially Poland, the Baltic, and the Benelux countries. In central Europe, soil moisture drought peaked in early February and March compared to a delayed peak in April in Poland and the Baltic and an even later peak in Ukraine and Moldova towards the end of 2019. The low flow situation in central Europe and Scandinavia partly recovered, although severe hazard levels were still detected from July to September. The eastern European countries showed an overall increase in low flow severity peaking earlier in the year (April and May). In addition, FAPAR was less severe in 2019 for most months and most of Europe, while south-eastern Europe and the Balkans showed increased hazard severity.

In southern and south-eastern Europe, the hydro-climatic conditions of 2018 and 2019 differed from the rest of Europe. In 2018, Spain, Portugal, and Italy had recovered from drought conditions, but deficits again developed in early 2019. In south-eastern Europe, winter 2018–2019 precipitation deficits were detected across much of the Balkan Peninsula, as well as in Slovakia. In Ukraine, Moldova, and Romania, the 2018 event was moderate in the second half of the year, and further rainfall deficits accumulated during winter, which led to rising soil moisture deficits from summer 2019 to the end of 2020.

3.2 The drought events of 2018 and 2019 – perception and management

The online survey yielded contributions by 712 respondents from 28 European countries (Fig. 2a) with the number of responses varying by country, i.e. from a single ex-




Figure 1. (a) Drought hazard conditions for 2018 and 2019 across the European continent according to the European Drought Observatory indicator fact sheets (<https://edo.jrc.ec.europa.eu/edov2/php/index.php?id=1101>, last access: 1 April 2022). Data are presented as the proportion of the country's total area under severe (or extreme) drought hazard conditions. Standardised Precipitation Index (SPI) is shown for accumulation periods of 3, 6, 9, and 12 months for August (AUG), September (SEP), October (OCT), and December (DEC). (b) SM is Soil Moisture Anomaly/Index, LFI is Low Flow Index, and fAPAR is Fraction of Photosynthetically Active Radiation, all presented for the day of the year in the corresponding colouring: 2018 as area and 2019 as lines.




Figure 2. Water usage across Europe: (a) number of survey participants by country, (b) most important water resource by country, and (c) second most important water resource by country. Results are based on a pan-European survey designed by the International Association of Hydrological Science (IAHS) – Panta Rhei “Drought in the Anthropocene” working group and conducted in 28 countries.

pert (Romania) to over 100 replies (Sweden). The majority of the respondents were employees at governmental institutions (74 %) at different administrative levels, with expertise related to water management, environment, meteorology, and agriculture. Furthermore, private and public companies (operators of public water supply systems, hydropower plants; 13 %), scientific institutions (4 %), and other non-governmental organisations with a focus on environment and ecology (3 %) also contributed to the survey.

The importance of the water resources as perceived by the participants (under normal conditions) ranked differently across the continent (Fig. 2b and c). The participants were asked to rank a selection of water resources and were also able to add additional ones. The sources of “artificial recharge” were not specified. If the nationally averaged importance of water resources were ranked equally (e.g. regulated and individual groundwater use both ranked as second most important), their importance as rank no. 2 and no. 3 were also evaluated. Overall, the majority of the respondents selected groundwater as the most important resource ($\sim 35\%$ of all participants), followed by surface water from rivers (22 %), reservoirs (13 %), individual wells (11 %), and artificial groundwater recharge (11 %). Further “Other” water resources such as rainfall collectors, ponds, or water transfer systems were listed a few times ($< 1\%$). Specific spatial patterns of water resource importance were not apparent, although individual wells appeared to be more important in eastern Europe, and artificial groundwater recharge was highlighted in Italy, Hungary, and Bulgaria. In the case of Spain, the questionnaire was adapted to national specificities and resulted in fewer water-usage categories; here “regulated surface water” falls in the category of “surface water from reservoirs”. Accordingly water resources ranks were no. 1 regulated surface water and no. 2 groundwater. A more detailed national breakdown of Fig. 2 can be found in the Supplement (Fig. S2).

The use of a drought definition to categorise drought hazard varied markedly across Europe (Fig. 3). About 40 % of all participants did not have an operational drought definition in their public and private organisations, and a further 15 % did not know whether there was one. In contrast, for the Czech Republic, Spain, Italy, and France all participants had an operational drought reporting system. With regard to the participants’ affiliation (see Table S2), about 60 % of those working for governmental authorities did not have – or were not aware of – an operational drought definition, in contrast to private companies in which around 30 % were unaware of a drought definition. Overall, about 20 % defined drought by a single drought type index (such as meteorological drought), 15 % used two, and 10 % used three different drought indices. The majority of participants used meteorological and hydrological indices (30 % each), and about 15 % relied on soil moisture and vegetation conditions. Furthermore, drought impact information, such as vegetation activity (e.g. Normalised Difference Vegetation Index, NDVI), crop yields, and forest fire indices, was used, in addition to media reports. In Spain, the “Special Drought Management Plans” define two types of drought-related events: prolonged drought (meteorological), quantified by precipitation deficit over different time periods, and conjunctural water scarcity, identified through the assessment of available water resources. This question was not asked in Sweden and Poland, and in Latvia, drought definitions were not operationalised.

Following the drought definition question, respondents were asked whether an established governmental drought declaration system existed or if the declaration of drought situations was based on case-specific decisions (Fig. S2). An operational declaration scheme is defined here as an official government-implemented method of defining a drought situation, often including drought severity thresholds and pre-defined measures. Operational drought declaration schemes



Figure 3. Major categories of drought indices used across Europe as a fraction of total replies per country (number of replies in parentheses, total replies = 536). The mean index in each category (meteorology, soil moisture, hydrology, and vegetation) is weighted by the number of participants. Countries on the y axis are sorted according to their mean index value, i.e. the highest for the Czech Republic, lowest for Slovakia. The category Others ($n = 37$) comprises countries with fewer than 10 replies, namely Austria (9), Italy (8), Belgium (6), Latvia (6), Iceland (4), Denmark (3), and Romania (1). Replies from Sweden and Poland are not considered here as indices were not rated in these countries. Note that participants have different roles in their countries and thus might judge drought indices differently. Results are based on a pan-European survey designed by the IAHS – Panta Rhei “Drought in the Anthropocene” working group and conducted in 28 countries.

(at country or county level) were scarce across the continent, though these were found to be present in Spain, France, the Netherlands, and the Czech Republic. In the Czech Republic, drought declaration is based on the open national drought monitoring platform Intersucho (Trnka et al., 2020). The same platform is shared also by the Slovak Hydrometeorological Institute (Labudová et al., 2018).

In Spain, governmental drought declaration schemes are included in the Special Drought Management Plans approved at the river basin level, where each basin has adapted it to their specific context and characteristics. Outside of Spain, individual decisions on drought declarations are more commonly present in regions with a lack of fixed drought declaration schemes. In some countries, drought situations are declared by the “Emergency situations commission meetings”, for example in Lithuania, the Netherlands, and the UK, where a national water management centre and drought committee advise the government. In Latvia, Estonia, Austria, Bulgaria, and Denmark, more than half of the respondents

did not know of the existence of any governmental drought declaration scheme or were not sure that one existed.

In over 25 countries, the majority of participants (> 50%) responded positively to the question of whether future climate change may affect water resources (Fig. 4a). The majority of respondents expected the occurrence of droughts to “increase” or “strongly increase” in the (near-)future (Fig. 4b). However, no relationship could be established between the expected future changes in drought hazard and the degree to which climate change is considered in policies. In addition, the responses about the expectation of “the need for more regulation of water distribution to fewer consumers due to shortages in the future” were linked to neither the expectation of future drought occurrence nor climate change (Fig. 4c). For example, in the UK, around 15% of respondents agreed that more regulation will be needed with a majority expecting an increase in drought occurrence, whereas in North Macedonia, about 85% agreed on a need for increased future regulation, having a similar share in future




Figure 4. Perception of climate change effect on drought management in Europe shown as percentage of participants responding to questions about (a) whether future climate change will may affect water resources, (b) how droughts may change in future, and (c) whether drinking water providers in the future would have to distribute water to fewer consumers due to shortages, e.g. “rota cuts”. Results are based on a pan-European survey designed by the IAHS – Panta Rhei “Drought in the Anthropocene” working group and conducted in 28 countries.




Figure 5. Perception of drought risk management across Europe shown as percentage of participants in pie charts: (a) distribution of drought risk management plans and emergency action plans by country; (b) reasons for an absence of drought risk management by country and totals of selected reasons. Results are based on a pan-European survey designed by the IAHS – Panta Rhei “Drought in the Anthropocene” working group and conducted in 28 countries.

drought occurrence. Nevertheless, the need for more regulation is perceived to be less important by participants from northern European countries compared to those from the rest of Europe. As mentioned by the participants, future regulation is expected to take the form of an EU drought directive, ranking priorities, re-allocating water permits, technological enhancements to save water, water pricing, and general water usage restrictions.

Few participants indicated that their countries had drought management action plans (~ 10 %), although emergency action plans were more common (~ 25 %), and both plans were more common in western Europe compared to eastern Europe (Fig. 5a). The UK, the Netherlands, Belgium, Spain, Switzerland, Italy, and Montenegro were comparatively well prepared in this regard (> 75 % of the participants had an emergency action or management plan). The countries of




Figure 6. Median perception of drought severity and impacts in 2018 and 2019 across Europe. Sweden participated only in 2018. Results are based on a pan-European survey designed by the IAHS – Panta Rhei “Drought in the Anthropocene” working group and conducted in 28 countries.

Spain, Italy, and the Czech Republic had a high share of participants indicating drought risk management plans in operational mode. More than 150 participants in the whole survey indicated the intention to introduce new (or update existing) drought management plans. As indicated by the participants, management tools include strategies that range from the increasing water storage capacity or adapting farming practices to the development of legally binding drought risk management actions.

To better understand the reasons for an absence of drought management plans, participants were asked for a possible explanation, and answers were provided by the national experts as either pre-defined or free text options. At the country scale, “insufficient resources” and the perception that “drought is not seen as a risk” were the most frequent answers (Fig. 5b). For northern Europe and Austria, drought not being seen as a risk was highlighted most often, whereas for the eastern European countries, the “lack of legal obligation – no European drought directive” and “waiting for governmental advice” were selected by about 15 % of all participants. Further, a “lack of knowledge” (on drought risk) was more prevalent in western Europe, whereas a “lack of resources” (finance and capacity) were prominent in central and south-eastern Europe. Political issues (e.g. “waiting for governmental advice”, “lack of forcing – no EU drought directive”, and “political lack of knowledge”) were especially present in central

and eastern Europe but were less prominent in northern Europe.

With regard to communication and interaction during drought events, participants were asked whom they collaborate with to manage droughts (Fig. S3). On average, more than half of the participants collaborated with “other authorities” (e.g. county administrative boards or water authorities). About 45 % interacted with “other departments or companies within the municipality” and about 20 % with “land owners and independent experts (such as universities)”. About 20 % did not know about any collaboration, and 5 % of participants stated no existing collaboration.

3.3 Survey-based perception and management of the 2018 and 2019 droughts in Europe

The perception of the 2018 and 2019 drought events by the survey respondents showed country- and event-specific differences (Fig. 6). The participants could rank the hydroclimatic situation from extremely dry to wet; 2018 was mostly perceived as being “drier than normal”, and central and northern European countries in particular were perceived as having “very dry” conditions (Fig. 6a), with high proportions (25 %) of “extremely dry” conditions in the Czech Republic, Germany, Lithuania, North Macedonia, and Norway (Fig. S4). South-western European coun-

tries also perceived 2018 as being “drier than normal”. In Iceland, 2018 was perceived as being a wet year, in contrast to the dry conditions of 2019 which were perceived to be “one of the worst droughts” on record. For the rest of Europe, the 2019 drought was perceived as being less severe than 2018 with the exception of France and Ukraine with high variations between countries (Fig. 6b). The centre of the 2018 drought event shifted by the end of the year/beginning of 2019 from central and northern Europe towards the east. Wetter conditions in northern Europe translated into perceptions of no or less severe drought in Scandinavia and the Baltic, respectively. The hydro-climatic situation in 2019 was still perceived as being “very dry” (50 %) in France, Belgium, Germany, Slovakia, and Ukraine.

Drought management preparation in 2018 showed an east-to-west gradient; i.e. eastern, northern, and central European countries felt overall more “prepared”, while countries in western Europe perceived they were “not well” prepared (Fig. S5). The management of the 2018 drought event was generally perceived as being worse compared to 2019, except in some central and northern European countries. Most respondents thought that they were better prepared in 2019 due to the previous event that likely contributed to an earlier activation of emergency plans, if any. However, the perception of drought impacts only shows minor differences between the two drought events (Fig. 6c and d), with the exception of northern Europe. The Mediterranean and the Balkan countries perceived drought impacts as not severe or without impacts (e.g. Croatia) in both years, with a tendency towards a higher severity in 2019 for Black Sea countries. In central Europe, participants perceived that they were severely affected in 2018, and this perception extended towards eastern Europe in 2019. Scandinavia and the Baltic were only slightly affected in 2018 with a lower perceived severity in 2019. For the majority of respondents, the drought of 2018 played a crucial role in the perceived impacts of the 2019 drought event (Fig. S5). Most respondents perceived particularly negative consequences for agriculture, livestock farming, forestry, and public water supply in 2018 and 2019 compared to relatively minor consequences regarding air pollution and conflicts (Fig. S6). At a first glance, the perception for these sectors differs only slightly between the events. However, soil moisture impacts, such as agricultural losses, impacts on freshwater aquaculture and fisheries, or forest fires, were reported less frequently in 2019. Denmark, Norway, and the UK had substantially fewer perceived impacts in the 2019 event than 2018 event. Slightly more impacts were reported in 2019 for livestock farming in Ukraine, for forestry and terrestrial ecosystems in Belgium and Ukraine, for air quality in Bulgaria, Slovakia, North Macedonia, and Ukraine, and for water quality in Austria, the Czech Republic, and North Macedonia. In contrast, Iceland was only affected in 2019 with strong effects on agriculture and water quality.

4 Discussion

This is the first study that quantifies drought perception by water-management-related stakeholders at continental scale based on a participatory survey. The survey analysis shows high diversity in perceived drought impacts (Fig. S6), which reflects Europe’s pedo-climatic and socio-economic heterogeneity, as is also shown by Stahl et al. (2016). The monitored and perceived drought hazard differed in some places as a result of the different drivers of drought impacts: hazard, exposure, and vulnerability (IPCC, 2014; UNDRR, 2019). The diversity of impacted categories has been reported previously for similar drought events, e.g. in 1975, 1976, and 2003 (Stahl et al., 2016). Our findings corroborate those of Stahl et al. (2016), with different countries across Europe being affected by the hazard very differently. Large-scale weather patterns and differences in land surface properties play a crucial role in explaining this heterogeneity. For instance, Atlantic meridional dipole circulation anomalies have been found to be associated with northern European droughts as represented by the SPI-6 and SPEI-6 (Standardised Precipitation Evapotranspiration Index) indices (Kingston et al., 2015). The Scandinavian teleconnection pattern, which was unusually high in May and July 2018, resembles the large-scale atmospheric circulation pattern most associated with summer low flow in southern and eastern Scandinavia (Bakke et al., 2020). It should be also noted that the frequency of drought-related circulation patterns has been changing since the end of the 19th century with increasing frequencies over central Europe (e.g. Lhotka et al., 2020; Trnka et al., 2009). The unique conditions of Iceland, where major drought events cannot be compared to the rest of Europe, were also shown by Spinoni et al. (2015), most likely attributed to its location influenced by warm humid winds and enhanced by the Gulf Stream clashing with the cold Arctic winds from the north that generate frequent precipitation events (de Niet et al., 2020). Nevertheless, severe land degradation in Iceland has decreased the water holding capacity, making the land susceptible to hydrological droughts (Finger et al., 2016; Keesstra et al., 2018). Furthermore, Spinoni et al. (2019) showed that major drought events as indicated by SPI and SPEI in central and northern Europe, north-eastern Europe, and southern Europe do not occur simultaneously, which was also evident in our results focusing on 2018 and 2019. The multi-year drought character of 2018 and 2019 became evident when focusing on the monitored hydrological drought conditions in Belgium, Switzerland, the Czech Republic, Germany, Finland, Latvia, Luxembourg, Lithuania, Poland, and Sweden.

In general, the hazard severity perceived by the surveyed stakeholders corresponded well with the hazard severity monitored by the EDO, though with some exceptions. For example, in 2018, large areas of Sweden and Finland were affected by severe (or extreme) hazard conditions according to the EDO, but the hydro-climatological situation was per-




Figure 7. National representatives' joint opinion on (a) the actual state of drought management in their country, (b) the existence of a country-wide drought management plan, (c) the existence of national recommendations for actions in order to minimise drought risk, and (d) the benefit of an EU drought directive for their country, ordered by score (very good = 3; not good = 0).

ceived as being “dry”. In 2019, EDO reported a severe (or extreme) meteorological drought in Iceland and a severe (or extreme) soil moisture drought in the Baltic, but in both cases, the hazard severity was perceived as being less severe and indexed as being “drier than normal” to “dry”. In contrast, Norway’s participants perceived very dry conditions in 2018, but the proportion of monitored severe (or extreme) hazard conditions was low. These discrepancies could be attributed to the low awareness of the stakeholders for the drought conditions at a larger scale or the impact across different sectors or the discrepancy between impact indicator and affected sector. A contributing factor could also be issues with the standardised drought indices effectively characterising the drought conditions in some regions. In higher-latitude countries, a strong negative rainfall anomaly does not necessarily imply a deficit in water availability for, for example, plant water uptake or public water supply as storage is usually replenished after the snowmelt period (Cammalleri et al., 2016). As such, meteorological drought indices may not be appropriate to predict impacts and consequences for the management of hydrological or agricultural (soil moisture) droughts. The wide range of drought definitions and associated high number of operational declaration schemes – highlight the many obstacles when dealing with the complex inter- and transdisciplinary nature of drought impacts. A unique definition of drought that is valid across all regions and sectors is not possible in practice (Lloyd-Hughes, 2014), especially if sectors, such as agriculture and water supply, are based on different laws and managed by different authorities. An effective implementation of macro-regional drought risk management requires a more holistic interdisciplinary view. Thus, drought cannot be declared by a single index only; the entire water cycle has to be considered as droughts in different parts of the water cycle can lead to different impacts. Such a holistic view should start with initial meteorological drought (e.g. lower than normal precipitation often combined with higher than normal evaporation) causing a deficit in soil moisture, and if sustained for a sufficient time, it may manifest itself as a hydrological drought (i.e. a deficit in streamflow and groundwater).

Our pan-European survey reflects the opinions of water professionals belonging to mostly the public sector and pub-

licly owned companies. The perspectives of other citizens, local stakeholders, private companies, and non-governmental organisations were less well represented. Nevertheless, the fraction of respondents’ affiliations differs among countries and could thus have had an influence on the hereingeneralised portrayal of drought risk. A statistical relation between affiliation and “other” replies (such as drought management or reason for a lack of drought risk management) could not be found. Furthermore, sectoral and regional perceptions of drought risk might differ. For example, a hydropower production survey in southern Germany showed that legislative drought risk regulation is not desired by reservoir operators, who would nevertheless support the development of drought risk management coupled with integrated river basin management (Siebert et al., 2021).

The preferential use of meteorological and hydrological indices to define drought by the participants was found to be similar to the findings of Bachmair et al. (2016). The absence of dedicated drought risk management strategies in many European countries is evident (Fig. 5a) due to diverse and, in some cases, contradictory reasons (Fig. 5b). The country representatives were asked some broad questions on the state of national drought management and the potential for a European drought directive; the responses revealed an unsatisfactory state of national drought risk management in Europe (Fig. 7).

The existence of drought risk management plans or strategies tended to be higher in countries with more common water scarcity issues and more frequent drought events, such as those in the Mediterranean region (Tramblay et al., 2020). Moreover, only Spain’s Special Drought Management Plans (updated in 2018; Hervás-Gómez and Delgado-Ramos, 2019) were considered as comprehensive and sufficient by the Spanish national representatives. In addition, recent drought events may have forced governments to foster drought research and policy implementation, suggesting that a “memory of recent disasters” improves disaster management and potentially mitigates drought impacts (Di Baldassarre et al., 2013; Kreibich et al., 2017). Urquijo et al. (2016) stated that drought management is a combination of the history of water management and the frequency of drought, which is supported by our results from the Mediterranean

countries and the Netherlands. Furthermore, case-specific effects of drought may also drive the need for risk management. In the Netherlands, for example, hydrological drought can increase salt water intrusion, land subsidence, and the structural instability of dikes. The resulting damage of these hydrological drought impacts decreases water security in the long term, especially with regard to compound events. The engagement of non-governmental scientific groups also fosters drought risk management and particularly public and government awareness (e.g. the Czech Republic).

The diversity of drought management approaches reflects the diversity of Europe's hydro-climatic conditions and governance contexts. However, droughts do not respect national borders, and Europe has several shared river basins. In addition, climate change is estimated to increase drought severity and frequency globally and in Europe (Erian et al., 2021; Spinoni et al., 2018).

The majority of the survey participants and all national representatives agreed that a pan-European drought management approach would support national and cross-boundary drought preparedness both now and in the future. While collaborations between water managers and agencies within countries are at least partly in place, as indicated in the survey, the difference between preparedness and proactive approaches to lower drought risk in Europe varies widely. Participants mainly in central and southern Europe indicated “insufficient resources”, “lack of forcing”, and “waiting for governmental advice” as reasons for not having a drought risk management plan. Across all national representatives and a majority of survey participants, there was a consensus that an EU directive on drought risk management would be beneficial (whether or not countries are EU member states). Similar to the Floods Directive (EU, 2018), a common strategy should only set a coarse framework, delegating specific actions to the member states and especially regulating trans-boundary water management during drought. An EU directive would be especially beneficial in countries where water resource management governance is not centralised, with wide procedural discrepancies among the different administrative regions and basin authorities. Recently, the Global Assessment Report on Drought (Erian et al., 2021) highlighted that adaptive risk management and governance strategies are required as responses to complex risks such as drought by means of actions, processes, and institutions. A drought directive, following the example of the European Floods Directive, would force member states and candidate countries to act and encourage cooperation across borders addressing the regional scale of drought hazard, secure resources, and funding for drought risk research and most importantly initiate a common strategy to increase drought resilience. However, not all respondents of the survey fully shared this view, the main reason being that a pan-European approach would not be able to consider local specificities such as catchment physical characteristics, water infrastructure, water uses, and specific biodiversity needs. Accordingly, a pan-European ap-

proach should also be tailor-made such that it can be trusted by the users. Therefore specific indicators and actions can be tailored to local situations and needs, but a general framework should be guiding the application of these. Thus, common action (e.g. a drought risk management strategy) may be conducted at a very general, broad, and political level. At the operational or local level, clear and common guidelines may be needed, and the challenge is to be flexible enough to cover context-specific situations.

5 Conclusions

The pan-European survey on drought perception and management highlighted the heterogeneity in the perception of drought hazard, impacts, and management across the European continent. The reflection on the drought events in the 2018–2019 period illustrated Europe's vulnerability to drought and the variable state of preparedness to withstand drought in many countries. Even though the awareness of a future increase in drought risk is prevalent, drought is often still not considered as a risk in central, northern, and eastern Europe. Here, we showed that drought hazard perception matched the observed or monitored drought hazard. In contrast, the occurrence of drought impacts does not always follow the pattern of hazard severity and therefore requires assessment of drought beyond just the hazard. A relationship between national drought awareness and drought management strategies could not be established. Although a strong variability in drought risk management planning across the continent was evident, a common European strategy does not exist. As shown here, current national drought risk management practices range from a fundamental lack of legislation to country-wide operational drought risk management plans. Future research might expand this survey to further explore and highlight potential benefits of a European drought directive. To foster national resilience to drought, drought management should be included in national legislation.

The key message of this study is that macro-governmental guidance by the EU is believed to be beneficial for national and international drought risk management. Such guidance should set a general framework which allows for regional flexibility of management strategies. To foster this kind of progress, sector-specific databases on drought impacts, such as the EDII, are required to show and quantify the varied impacts of past droughts and increase public awareness in order to encourage political action. Going a step further, such information should be hosted by (inter-)national drought risk monitoring systems presenting sector-specific drought risk.

As the first major steps towards a more unified drought risk management in Europe, we recommend

1. the inclusion of a clear definition of drought in the Water Framework Directive, considering different types of drought, as well as their spatial and temporal occurrence,

2. the development of impact-driven, regional- and sector-specific guidance on drought indices, and
3. the formation of an inter- and transdisciplinary collaborative EU working group focusing on drought risk management and estimation of the potential benefits and downsides of a European Drought Directive.

Code availability. Data analysis code is not available. The analysis followed standard statistical routines and can be reproduced by the methodological explanations in the text.

Data availability. Drought monitoring and survey data can be accessed via <https://doi.org/10.34730/ae96ed78875c4caa9ee5c25c2e2f711a> (Blauhut, 2021).

Supplement. The supplement related to this article is available online at: <https://doi.org/10.5194/nhess-22-2201-2022-supplement>.

Author contributions. VB developed the basis of the study in cooperation with the major contributing authors, MS, LA, MIB, CT, and DEW. VB developed the network of national representatives, set up all questionnaires, and proceeded with all data analyses in cooperation with MS. LA, MIB, CT, and DEW participated in the formal analysis of the results and drafting of the original draft. CC provided drought hazard information from the JRC and participated in the formal analysis of the result. LJB did an intense job in revising spelling, grammar, and phrasing of multiple versions of the manuscript. All authors, a.k.a. national representatives, shaped the catalogue of questions, translated the questionnaire into the national language, developed a national network of contributors, distributed and quality checked the questionnaire, revised results, and contributed to the manuscript. Furthermore, all authors contributed to the questionnaire of national representatives.

Competing interests. At least one of the (co-)authors is a member of the editorial board of *Natural Hazards and Earth System Sciences*. The peer-review process was guided by an independent editor, and the authors also have no other competing interests to declare.

Disclaimer. Publisher's note: this study is based on the NUTS-level classification system (and GIS data) as provided by Eurostat for the year 2016. Accordingly, the pan-European survey followed the NUTS regions and countries as given.

Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Special issue statement. This article is part of the special issue "Drought vulnerability, risk, and impact assessments: bridging the science-policy gap". It is a result of the EGU General Assembly 2020, 3–8 May 2020.

Acknowledgements. This research was carried out within the interdisciplinary research project DRIeR. Special thanks go to the IAHS – Panta Rhei "Drought in the Anthropocene" working group who originally designed the study.

Financial support. The project is supported by the *Wassernetzwerk Baden-Württemberg* (Water Research Network of the State Baden-Württemberg), which is funded by the *Ministerium für Wissenschaft, Forschung und Kunst Baden-Württemberg* (Ministry of Science, Research and the Arts of the State Baden-Württemberg) (grant no. AZ. 7532.21/2.1.6) and Maa- ja vesitekniiikan tuki ry foundation. Doris E. Wendt acknowledges her support as part of the NERC-funded Groundwater Drought Initiative (NE/R004994/1). Lucy J. Barker was supported by the Natural Environment Research Council (NE/R016429/1) as part of the UK-SCAPE programme delivering National Capability. The contributions of Mirek Trnka, Lenka Bartošová, and Jaak Jaagus have been supported by SustES – Adaptation strategies for sustainable ecosystem services and food security under adverse environmental conditions (CZ.02.1.01/0.0/0.0/16_019/0000797).

This open-access publication was funded by the University of Freiburg.

Review statement. This paper was edited by Maria-Carmen Llasat and reviewed by two anonymous referees.

References

- Aldunce, P., Araya, D., Sapiain, R., Ramos, I., Lillo, G., Urquiza, A., and Garreaud, R.: Local perception of drought impacts in a changing climate: The mega-drought in central Chile, *Sustainability*, 9, 205, <https://doi.org/10.3390/su9112053>, 2017.
- Bachmair, S., Stahl, K., Collins, K., Hannaford, J., Acreman, M., Svoboda, M., Knutson, C., Smith, K. H., Wall, N., Fuchs, B., and Crossman, N. D.: Drought indicators revisited: the need for a wider consideration of environment and society, *Wiley Interdisciplin. Rev. Water*, 3, 516–536, 2016.
- Bakke, S. J., Ionita, M., and Tallaksen, L. M.: The 2018 northern European hydrological drought and its drivers in a historical perspective, *Hydrol. Earth Syst. Sci.*, 24, 5621–5653, <https://doi.org/10.5194/hess-24-5621-2020>, 2020.
- Barker, L. J., Hannaford, J., Parry, S., Smith, K. A., Tanguy, M., and Prudhomme, C.: Historic hydrological droughts 1891–2015: systematic characterisation for a diverse set of catchments across the UK, *Hydrol. Earth Syst. Sci.*, 23, 4583–4602, <https://doi.org/10.5194/hess-23-4583-2019>, 2019.
- Baruth, B., Bassu, S., Bussay, A., Ceglar, A., Cerrani, I., Chemin, Y., De Palma, P., Fumagalli, D., Lecerf, R., Manfron, G., Nisini Scacchiafichi, L., Panarello, L., Ronchetti, G., Seguni, L., Toreti,

- A., Van Den Berg, M., Zajac, Z., Zucchini, A., and Biavetti, I.: Crop monitoring in Europe, in: JRC MARS Bulletin, Vol. 28, No. 9, edited by: Van Den Berg, M., Baruth, B., Niemeier, S., Van Der Velde, M., and Manfron, G., Publications Office of the European Union, Luxembourg, JRC120746, <https://publications.jrc.ec.europa.eu/repository/handle/JRC120746> (last access: 1 April 2022), 2020.
- Beillouin, D., Schauburger, B., Bastos, A., Ciais, P., and Makowski, D.: Impact of extreme weather conditions on European crop production in 2018, *Philos. T. Roy. Soc. Lond. B*, 375, 20190510, <https://doi.org/10.1098/rstb.2019.0510>, 2020.
- Bastawesy, M. E.: Hydrological scenarios of the Renaissance Dam in Ethiopia and its hydro-environmental impact on the Nile downstream, *J. Hydrol. Eng.*, 20, 04014083, [https://doi.org/10.1061/\(ASCE\)HE.1943-5584.0001112](https://doi.org/10.1061/(ASCE)HE.1943-5584.0001112), 2015.
- Blauhut, V.: The triple complexity of drought risk analysis and its visualisation via mapping: a review across scales and sectors, *Earth-Sci. Rev.*, 210, 103345, <https://doi.org/10.1016/j.earscirev.2020.103345>, 2020.
- Blauhut, V.: European drought management and perception – insights from 2018 & 2019 events, B2SHARE [data set], <https://b2share.fz-juelich.de/records/ae96ed78875c4caa9ee5c25c2e2f711a>, last access: 11 February 2022), 2021.
- Blauhut, V., Stahl, K., and Kohn, I.: The dynamics of vulnerability to drought from an impact perspective, in: Drought: Research and Science-Policy Interfacing, edited by: Andreu, H. J., Solera, A., Paredes-Arquiola, J., Haro-Montegudo, D., and van Lanen (Ed.), CRC Press, London, 349–354, ISBN 9780429226380, 2015.
- Blauhut, V., Stahl, K., Stagger, J. H., Tallaksen, L. M., De Stefano, L., and Vogt, J.: Estimating drought risk across Europe from reported drought impacts, drought indices, and vulnerability factors, *Hydrol. Earth Syst. Sci.*, 20, 2779–2800, <https://doi.org/10.5194/hess-20-2779-2016>, 2016.
- Boergens, E., Güntner, A., Dobsław, H., and Dahle, C.: Quantifying the Central European Droughts in 2018 and 2019 with GRACE Follow-On, *Geophys. Res. Lett.*, 47, e2020GL087285, <https://doi.org/10.1029/2020GL087285>, 2020.
- Büntgen, U., Urban, O., Krusic, P. J., Rybníček, M., Kolář, T., Kyncl, T., Ač, A., Koňasová, E., Čáslavský, J., Esper, J., Wagner, S., and Trnka, M.: Recent European drought extremes beyond Common Era background variability, *Nat. Geosci.*, 14, 190–196, <https://doi.org/10.1038/s41561-021-00698-0>, 2021.
- Cammalleri, C., Naumann, G., Mentaschi, L., Formetta, G., Forzieri, G., Gosling, S., Bisselink, B., De Roo, A., and Feyen, L.: Global warming and drought impacts in the EU, Publications Office of the European Union, <https://doi.org/10.2760/597045>, 2020.
- Cammalleri, C., Micale, F., and Vogt, J.: A novel soil moisture-based drought severity index (DSI) combining water deficit magnitude and frequency, *Hydrol. Process.*, 30, 289–301, <https://doi.org/10.1002/hyp.10578>, 2016.
- Cheval, S., Busuioc, A., Dumitrescu, A., and Birsan, M.-V.: Spatiotemporal variability of meteorological drought in Romania using the standardized precipitation index (SPI), *Clim. Res.*, 60, 235–248, 2014.
- Cindrić Kalin, K., Prtenjak, M. T., Herceg-Bulić, I., Mihajlović, D., and Pasarić, Z.: Analysis of the extraordinary 2011/2012 drought in Croatia, *Theor. Appl. Climatol.*, 123, 503–522, 2016.
- Dai, A.: Increasing drought under global warming in observations and models, *Nat. Clim. Change*, 3, 52–58, 2013.
- de Brito, M. M.: Compound and cascading drought impacts do not happen by chance: A proposal to quantify their relationships, *Sci. Total Environ.*, 778, 146236, <https://doi.org/10.1016/j.scitotenv.2021.146236>, 2021.
- de Niet, J., Finger, D. C., Bring, A., Egilson, D., Gustafsson, D., and Kalantari, Z.: Benefits of Combining Satellite-Derived Snow Cover Data and Discharge Data to Calibrate a Glaciated Catchment in Sub-Arctic Iceland, *Water*, 12, 975, <https://doi.org/10.3390/w12040975>, 2020.
- Di Baldassarre, G., Viglione, A., Carr, G., Kuil, L., Salinas, J. L., and Blöschl, G.: Socio-hydrology: conceptualising human-flood interactions, *Hydrol. Earth Syst. Sci.*, 17, 3295–3303, <https://doi.org/10.5194/hess-17-3295-2013>, 2013.
- DriDanube-Watch: Drought 2018 – DriDanube watch, Vol. 37), <https://drive.google.com/file/d/1m-qobw41JGauZdIANlS1NPaRe8SQ2Gp9/view> (last access: 1 April 2021), 2018.
- Erian, W., Pulwarty, R., Vogt, J. V., AbuZeid, K., Bert, F., Bruntrup, M., El-Askary, H., de Estrada, M., Gaupp, F., Grundy, M., Hadwen, T., Hagenlocher, M., Kairu, G., Lamhauge, N., Li, W., Mahon, R., Maia, R., Martins, E. S. P. R., Meza, I., de los Milagos Skansi, M., Moderc, A., Naumann, G., Negri, R., Partex, S. P., Podesta, G., Quesada, M., Rakhmatova, Riley, J. E., Rudari, R., Shanmugasundaram, J., Silveira Reis, S., Singh, C., Spennemann, P., Srinivasan, G., Stefanski, R., Sušnik, A., Svoboda, M., Trotman, A., Tsegai, D., Unver, O., Van Meerbeeck, C., and Wens, M.: GAR Special Report on Drought 2021, UNDRR – United Nations Office for Disaster Risk Reduction, ISBN 9789212320274, <https://www.undrr.org/media/49386/download> (last access: 1 April 2022), 2021.
- EU: Directive 2007/60/EC of the European Parliament and of the council of 23 October 2007 on the assessment and management of flood risks, *Off. J. Eur. Union Legis.*, 50, 27–34, 2018.
- European Commission: A Blueprint to Safeguard Europe’s Water Resources, COM (2012) 673 final, Brussels, <http://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX:52012DC0673> (last access: 1 April 2022), 2012.
- European Commission: Fitness Check of the Water Framework Directive and the Floods Directive, 1–184, [https://ec.europa.eu/environment/water/fitness_check_of_the_eu_water_legislation/documents/WaterFitnessCheck-SWD\(2019\)439-web.pdf](https://ec.europa.eu/environment/water/fitness_check_of_the_eu_water_legislation/documents/WaterFitnessCheck-SWD(2019)439-web.pdf) (last access: 1 April 2022), 2019.
- Finger, D., Þórsson, J., Pétursdóttir, Þ., and Halldórsson, G.: Enhancing the resilience of water resources through land restoration in Rangárvellir, Iceland – an overview of the HydroResilience project, SER Europe Knowledge Base, Extended abstract 10th European Conference on Ecological Restoration, Freising, Germany, 5 pp., ISSN 2295-5704, 2016.
- Finger, D. C.: The value of satellite retrieved snow cover images to assess water resources and the theoretical hydropower potential in ungauged mountain catchments, *Jökull*, 68, 47–66, 2018.
- García-Herrera, R., Garrido-Perez, J. M., Barriopedro, D., Ordóñez, C., Vicente-Serrano, S. M., Nieto, R., Gimeno, L., Sorí, R., and

- Yiou, P.: The European 2016/17 drought, *J. Climate*, 32, 3169–3187, <https://doi.org/10.1175/JCLI-D-18-0331.1>, 2019.
- Hagenlocher, M., Meza, I., Anderson, C. C., Min, A., Renaud, F. G., Walz, Y., Siebert, S., and Sebesvari, Z.: Drought vulnerability and risk assessments: state of the art, persistent gaps, and research agenda, *Environ. Res. Lett.*, 14, 83002, <https://doi.org/10.1088/1748-9326/ab225d>, 2019.
- Hanger-Kopp, S. and Palka, M.: Exploring Drought Resilience Through a Drought Risk Management Lens in Austria, in: *Disaster Risk Reduction and Resilience, Disaster and Risk Research: GADRI Book Series*, edited by: Yokomatsu, M. and Hochrainer-Stigler, S., Springer, Singapore, https://doi.org/10.1007/978-981-15-4320-3_7, 2020.
- Hänsel, S., Ustrul, Z., Łupikasza, E., and Skalak, P.: Assessing seasonal drought variations and trends over Central Europe, *Water Resour.*, 127, 53–57, 2019.
- Hari, V., Rakovec, O., Markonis, Y., Hanel, M., and Kumar, R.: Increased future occurrences of the exceptional 2018–2019 Central European drought under global warming, *Scient. Rep.*, 10, 12207, <https://doi.org/10.1038/s41598-020-68872-9>, 2020.
- Helmert, J., Şensoy Şorman, A., Alvarado Montero, R., De Michele, C., De Rosnay, P., Dumont, M., Finger, D. C., Lange, M., Picard, G., Potopová, V., Pullen, S., and Vikhamar-Schuler, D.: An Review of Snow Data Assimilation Methods for Hydrological, Land Surface, Meteorological and Climate Models: Results from a COST HarmoSnow Survey, *Geosciences*, 8, 489, <https://doi.org/10.3390/geosciences8120489>, 2018.
- Hervás-Gámez, C. and Delgado-Ramos, F.: Drought management planning policy: From Europe to Spain, *Sustainability*, 11, 1–26, <https://doi.org/10.3390/su11071862>, 2019.
- Hisdal, H. and Tallaksen, L. M.: Estimation of regional meteorological and hydrological drought characteristics: a case study for Denmark, *J. Hydrol.*, 281, 230–247, [https://doi.org/10.1016/S0022-1694\(03\)00233-6](https://doi.org/10.1016/S0022-1694(03)00233-6), 2003.
- Howarth, W.: Going with the flow: Integrated Water Resources Management, the EU Water Framework Directive and ecological flows, *Legal Stud.*, 38, 298–319, <https://doi.org/10.1017/lst.2017.13>, 2018.
- Ionita, M., Tallaksen, L. M., Kingston, D. G., Stagge, J. H., Laaha, G., Van Lanen, H. A. J., Scholz, P., Chelcea, S. M., and Haslinger, K.: The European 2015 drought from a climatological perspective, *Hydrol. Earth Syst. Sci.*, 21, 1397–1419, <https://doi.org/10.5194/hess-21-1397-2017>, 2017.
- IPCC: *Climate Change 2014 Impact, Adaption, and Vulnerability Part A: Global and sectoral aspects*, New York, USA, <https://doi.org/10.1017/CBO9781107415324.004>, 2014.
- Jaagus, J., Aasa, A., Aniskevich, S., Boincean, B., Bojariu, R., Briede, A., Danilovich, I., Domínguez Castro, F., Dumitrescu, A., Labuda, M., and Labudová, L.: Long-term changes in drought indices in eastern and central Europe, *Int. J. Climatol.*, in preparation, 2021.
- Keesstra, S., Nunes, J., Novara, A., Finger, D., Avelar, D., Kalantari, Z., and Cerdà, A.: The superior effect of nature based solutions in land management for enhancing ecosystem services, *Sci. Total Environ.*, 610, 997–1009, <https://doi.org/10.1016/j.scitotenv.2017.08.077>, 2018.
- Kingston, D. G., Stagge, J. H., Tallaksen, L. M., and Hannah, D. M.: European-Scale Drought: Understanding Connections between Atmospheric Circulation and Meteorological Drought Indices, *J. Climate*, 28, 505–516, <https://doi.org/10.1175/JCLI-D-14-00001.1>, 2015.
- Kreibich, H., Di Baldassarre, G., Vorogushyn, S., Aerts, J. C. J. H., Apel, H., Aronica, G. T., Arnbjerg-Nielsen, K., Bouwer, L. M., Bubeck, P., Caloiero, T., Chinh, D. T., Cortès, M., Gain, A. K., Giampá, V., Kuhlicke, C., Kundzewicz, Z. W., Llasat, M. C., Mård, J., Matczak, P., Mazzoleni, M., Molinari, D., Dung, N. V., Petrucci, O., Schröter, K., Slager, K., Thielen, A. H., Ward, P. J., and Merz, B.: Adaptation to flood risk: Results of international paired flood event studies, *Earth's Future*, 5, 953–965, <https://doi.org/10.1002/2017EF000606>, 2017.
- Kreibich, H., Blauhut, V., Aerts, J. C. J. H., Bouwer, L. M., Van Lanen, H. A. J., Mejia, A., Mens, M., and Van Loon, A. F.: How to improve attribution of changes in drought and flood impacts, *Hydrolog. Sci. J.*, 64, 1–18, <https://doi.org/10.1080/02626667.2018.1558367>, 2019.
- Laaha, G., Gauster, T., Tallaksen, L. M., Vidal, J.-P., Stahl, K., Prudhomme, C., Heudorfer, B., Vlnas, R., Ionita, M., Van Lanen, H. A. J., Adler, M.-J., Caillouet, L., Delus, C., Fendekova, M., Gailliez, S., Hannaford, J., Kingston, D., Van Loon, A. F., Mediero, L., Osuch, M., Romanowicz, R., Sauquet, E., Stagge, J. H., and Wong, W. K.: The European 2015 drought from a hydrological perspective, *Hydrol. Earth Syst. Sci.*, 21, 3001–3024, <https://doi.org/10.5194/hess-21-3001-2017>, 2017.
- Labudová, L., Ivaňáková, G., Trnka, M., and Bartošová, L.: Monitoring of drought impacts and the DriDanube project (Tromp Foundation Travel Award), in: Vol. 15, *EMS Annual Meeting Abstracts*, 3–7 September 2018, Budapest, Hungary, EMS2018-34-1, 2018.
- Lhotka, O., Trnka, M., Kyselý, J., Markonis, Y., Balek, J., and Možný, M.: Atmospheric circulation as a factor contributing to increasing drought severity in central Europe, *J. Geophys. Res.-Atmos.*, 125, e2019JD032269, <https://doi.org/10.1029/2019JD032269>, 2020.
- Lloyd-Hughes, B.: The impracticality of a universal drought definition, *Theor. Appl. Climatol.*, 117, 607–611, <https://doi.org/10.1007/s00704-013-1025-7>, 2014.
- McKee, T. B., Doesken, N. J., and Kleist, J.: The relationship of drought frequency and duration to time scales, in: Vol. 17, *Proceedings of the 8th Conference on Applied Climatology*, 17–22 January 1993, Anaheim, California, 179–183, 1993.
- Moravec, V., Markonis, Y., Rakovec, O., Svoboda, M., Trnka, M., Kumar, R., and Hanel, M.: Europe under multi-year droughts: how severe was the 2014–2018 drought period?, *Environ. Res. Lett.*, 16, 034062, <https://doi.org/10.1088/1748-9326/abe828>, 2021.
- Radeva, K., Nikolova, N., and Gera, M.: Assessment of hydro-meteorological drought in the Danube Plain, Bulgaria, *Hrvatski Geografski Glasnik*, 80, 7–25, 2018.
- Rita, A., Camarero, J. J., Nolè, A., Borghetti, M., Brunetti, M., Pergola, N., Serio, C., Vicente-Serrano, S. M., Tramutoli, V., and Ripullone, F.: The impact of drought spells on forests depends on site conditions: The case of 2017 summer heat wave in southern Europe, *Global Change Biol.*, 26, 851–863, 2020.
- Rosner, B., Benedict, I., van Heerwaarden, C., Weerts, A., Hazeleger, W., Bissolli, P., and Trachte, K.: The long heat wave and drought in Europe in 2018, *B. Am. Meteorol. Soc.*, 100, S222–S223, 2019.

- Salmoral, G., Ababio, B., and Holman, I. P.: Drought impacts, coping responses and adaptation in the UK outdoor livestock sector: Insights to increase drought resilience, *Land*, 9, 202, <https://doi.org/10.3390/LAND9060202>, 2020.
- Schuldts, B., Buras, A., Arend, M., Vitasse, Y., Beierkuhnlein, C., Damm, M., Gharun, M., Grams, T. E. E., Hauck, M., Hajek, P., Hartmann, H., Hiltbrunner, H., Hoch, G., Holloway-Phillips, M. H., Körner, C., Larysch, E., Lübke, T., Nelson, D. B., Rammig, A., Rigling, A., Rose, L., Ruehr, N. K., Schumann, K., Weiser, F., Werner, C., Wohlgenuth, T., Zang, C. S. and Kahmen, A.: A first assessment of the impact of the extreme 2018 summer drought on Central European forests, *Basic Appl. Ecol.*, 45, 86–103, <https://doi.org/10.1016/j.baae.2020.04.003>, 2020.
- Siebert, C., Blauhut, V., and Stahl, K.: Drought and its implications for the public water supply sector in Baden-Württemberg, Germany: impacts, management, perception, *Wasser-Wirtschaft* 6/2021, <https://www.springer.com/journal/35147> (last access: 1 April 2022), 2021.
- Spinoni, J., Naumann, G., Vogt, J. V., and Barbosa, P.: The biggest drought events in Europe from 1950 to 2012, *J. Hydrol.: Reg. Stud.*, 3, 509–524, <https://doi.org/10.1016/j.ejrh.2015.01.001>, 2015.
- Spinoni, J., Vogt, J. V., Naumann, G., Barbosa, P., and Dosio, A.: Will drought events become more frequent and severe in Europe?, *Int. J. Climatol.*, 38, 1718–1736, <https://doi.org/10.1002/joc.5291>, 2018.
- Spinoni, J., Barbosa, P., De Jager, A., McCormick, N., Naumann, G., Vogt, J. V., Magni, D., Masante, D., and Mazzeschi, M.: A new global database of meteorological drought events from 1951 to 2016, *J. Hydrol.: Reg. Stud.*, 22, 100593, <https://doi.org/10.1016/j.ejrh.2019.100593>, 2019.
- Stahl, K., Kohn, I., Blauhut, V., Urquijo, J., De Stefano, L., Acácio, V., Dias, S., Stage, J. H., Tallaksen, L. M., Kampragou, E., Van Loon, A. F., Barker, L. J., Melsen, L. A., Bifulco, C., Musolino, D., de Carli, A., Massarutto, A., Assimakopoulos, D., and Van Lanen, H. A. J.: Impacts of European drought events: insights from an international database of text-based reports, *Nat. Hazards Earth Syst. Sci.*, 16, 801–819, <https://doi.org/10.5194/nhess-16-801-2016>, 2016.
- Stein, U., Özerol, G., Tröltzsch, J., Landgrebe, R., Szendrenyi, A., and Vidaurre, R.: European Drought and Water Scarcity Policies, in: *Governance for Drought Resilience: Land and Water Drought Management in Europe*, edited by: Bressers, H., Bressers, N., and Larrue, C., Springer International Publishing, Cham, 17–43, https://doi.org/10.1007/978-3-319-29671-5_2, 2016.
- Teutschbein, C., Lundkvist, E., Albrecht, F., and Blicharska, M.: A culture of proactive drought management? Unraveling the perception and management of droughts in Swedish municipalities, *Geophys. Res. Abstr.*, 21, 1, 2019.
- Thompson, R. L., Broquet, G., Gerbig, C., Koch, T., Lang, M., Monteil, G., Munassar, S., Nickless, A., Scholze, M., Ramonet, M., Karstens, U., van Schaik, E., Wu, Z., and Rödenbeck, C.: Changes in net ecosystem exchange over Europe during the 2018 drought based on atmospheric observations, *Philos. T. Roy. Soc. Lond. B*, 375, 20190512, <https://doi.org/10.1098/rstb.2019.0512>, 2020.
- Tramblay, Y., Koutroulis, A., Samaniego, L., Vicente-Serrano, S. M., Volaire, F., Boone, A., Le Page, M., Llasat, M. C., Albergel, C., Burak, S., Cailleret, M., Cindric, K., Davi, H., Dupuy, J., Greve, P., Grillakis, M., Hanich, L., Jarlan, L., Martin-Stpaul, N., and Polcher, J.: Challenges for Drought Assessment in the Mediterranean Region under Future Climate Scenarios, *Earth-Sci. Rev.*, 210, 103348, <https://doi.org/10.1016/j.earscirev.2020.103348>, 2020.
- Trnka, M., Kyselý, J., Možný, M., and Dubrovský, M.: Changes in Central-European soil-moisture availability and circulation patterns in 1881–2005, *Int. J. Climatol.*, 29, 655–672, <https://doi.org/10.1002/joc.1703>, 2009.
- Trnka, M., Hayes, M., Jurečka, F., Anderson, M., Brázdil, R., Brown, J., and Feng, S.: Priority questions in multidisciplinary drought research, *Clim. Res.*, 75, 241–260, <https://doi.org/10.3354/cr01509>, 2018.
- Trnka, M., Hlavinka, P., Možný, M., Semerádová, D., Štěpánek, P., Balek, J., Bartošová, L., Zahradníček, P., Bláhová, M., Skalák, P., and Farda, A.: Czech Drought Monitor System for monitoring and forecasting agricultural drought and drought impacts. *Int. J. Climatol.*, 40, 5941–5958, <https://doi.org/10.1002/joc.6557>, 2020.
- Turner, S., Barker, L. J., Hannaford, J., Muchan, K., Parry, S., and Sefton, C.: The 2018/2019 drought in the UK: a hydrological appraisal, *Weather*, 78, 248–253, <https://doi.org/10.1002/wea.4003>, 2021.
- UNCCD: Drought Resilience, Adaptation and Management Policy Framework: Supporting Technical Guidelines, 17 pp., <https://www.unccd.int/resources/manuals-and-guides/drought-resilience-adaptation-and-management-policy-framework> (last access: 1 April 2022), 2019.
- UNDRR – United Nations Office for Disaster Risk Reduction: Global Assessment Report on Disaster Risk Reduction, United Nations Office for Disaster Risk Reduction, Geneva, Switzerland, <https://gar.undrr.org/> (last access: 1 April 2022), 2019.
- Urquijo, J., Pereira, D., Dias, S., and De Stefano, L.: A methodology to assess drought management as applied to six European case studies, *Int. J. Water Resour. Dev.*, 627, 1–24, <https://doi.org/10.1080/07900627.2016.1174106>, 2016.
- Van Lanen, H. A. J., Laaha, G., Kingston, D. G., Gauster, T., Ionita, M., Vidal, J.-P., Vlnas, R., Tallaksen, L. M., Stahl, K., Hannaford, J., Delus, C., Fendekova, M., Mediero, L., Prudhomme, C., Rets, E., Romanowicz, R. J., Gailliez, S., Wong, W. K., Adler, M.-J., Blauhut, V., Caillouet, L., Chelcea, S., Frolova, N., Gudmundsson, L., Hanel, M., Haslinger, K., Kireeva, M., Osuch, M., Sauquet, E., Stage, J. H., and Van Loon, A. F.: Hydrology needed to manage droughts: the 2015 European case, *Hydrol. Process.*, 30, 3097–3104, <https://doi.org/10.1002/hyp.10838>, 2016.
- Vuković, M., Voza, D., Štrbac, N., and Takić, L.: Cooperation over International Water Resources: a Case from the Danube River Basin, *Sociológia*, 46, 320–342, 2014.
- Zoth, P., Caillet, V., and Mager, U.: Herausforderung und Realität eines Dürremanagements in Baden-Württemberg: ein Bericht aus dem Forschungsnetzwerk DRiEr, VBIB – Verwaltungsblätter Baden-Württemberg, 133–142, ISSN 0720-2407, 2019.